


OXYFAST (OXYCODONE) 20MG/1ML

Instruction Sheet

Oxyfast is indicated to control pain or relieve breathing difficulties.

Directions: Use 1 ml syringe or the eyedropper that comes with the medication.

- To give 5mg draw up .25ml
- To give 10mg draw up .5ml
- To give 15mg draw up .75ml
- To give 20mg draw up 1 ml (ml's are the same as cc's)



Please use extra care when preparing the medication. It is important to use the correct amount. In the home you can reuse the syringes and eyedropper after rinsing with soap, warm water and air drying. Prior to washing, take the syringe apart.

Oxyfast (Oxycodone) may be swallowed, however if the patient is unable to swallow or has difficulty swallowing, you may place the medication under the tongue or in the cheek between the teeth and cheek. Oxyfast (Oxycodone) will then be absorbed into the oral mucosa tissue.

Oxyfast (Oxycodone) is a rapid acting medication and is usually effective within ten to twenty minutes.

Side effects of Oxyfast (Oxycodone) include drowsiness, dizziness, lightheadedness, itching, mental changes. Do not drive a vehicle, operate machinery, or drink alcohol when taking this medication. Do not share this medication with anyone else. Please notify your nurse if you are experiencing any side effects of the medication.



Hospice Alliance™

Ask for us by name

Your compassionate, community non-profit hospice, incorporated in 1981.

OXYFAST (OXYCODONE) 20MG/1ML

Instruction Sheet (continued)

Taking opioid medications may cause constipation. Please discuss a bowel program with your nurse to ensure that constipation is not an issue.

If you have any questions relating to this, or any medication, please do not hesitate to call us; day or night at 262-652-4400 or 800-830-8344.